

Zero Energy Bound States in Three-Particle Systems

Dmitry K. Gridnev

*FIAS, Ruth-Moufang-Straße 1, D-60438 Frankfurt am Main, Germany**

Abstract

We consider a 3-body system in $\mathbb{R}^3$ with non-positive potentials and non-negative essential spectrum. Under certain requirements on the fall off of pair potentials it is proved that if at least one pair of particles has a zero energy resonance then a square integrable zero energy ground state of three particles does not exist. This complements the analysis in [1], where it was demonstrated that square integrable zero energy ground states are possible given that in all two-body subsystems there is no negative energy bound states and no zero energy resonances. As a corollary it is proved that one can tune the coupling constants of pair potentials so that for any given $R, \epsilon > 0$: (a) the bottom of the essential spectrum is at zero; (b) there is a negative energy ground state $\psi(\xi)$, where $\int |\psi(\xi)|^2 = 1$; (c) $\int_{|\xi| \leq R} |\psi(\xi)|^2 < \epsilon$.

*On leave from: Institute of Physics, St. Petersburg State University, Ulyanovskaya 1, 198504 Russia

I. INTRODUCTION

In [1] under certain restrictions on pair potentials it was proved that the 3-body system, which is at the 3-body coupling constant threshold, has a square integrable state at zero energy if none of the 2-body subsystems is bound or has a zero energy resonance. Naturally, a question can be raised whether the condition on the absence of 2-body zero energy resonances is essential. Here we show that indeed it is. Namely, in this paper we prove that the 3-body ground state at zero energy can only be a resonance and not a L^2 state if at least one pair of particles has a resonance at zero energy. The method of proof is inspired by [2, 3]. In the last section we demonstrate that there do exist 3-body systems, where (a) each 2-body subsystem is unbound; (b) one 2-body subsystem is at the coupling constant threshold; (c) the 3-body system has a resonance at zero energy and bound states with the energy less or equal to zero. System like this can be constructed through appropriate tuning of the coupling constants.

II. A ZERO ENERGY RESONANCE IN A 2-BODY SYSTEM

In this section we shall use the method of [2] to prove a result similar to Lemma 2.2 in [3]. Let us consider the Hamiltonian of 2 particles in $\mathbb{R}^3$

$$h_{12}(\varepsilon) := -\Delta_x - (1 + \varepsilon)V_{12}(\alpha x), \quad (1)$$

where $\varepsilon \geq 0$ is a parameter and $\alpha = \hbar/\sqrt{2\mu_{12}}$ is a constant, which depends on the reduced mass μ_{12} [1]. Additionally, we require

R1 $0 \leq V_{12}(\alpha x) \leq b_1 e^{-b_2|x|}$, where $b_{1,2} > 0$ are constants.

R2 $h_{12}(0) \geq 0$ and $\sigma_{pp}(h_{12}(\varepsilon)) \cap (-\infty, 0) \neq \emptyset$ for $\varepsilon > 0$.

The requirement R2 means that $h_{12}(0)$ has a resonance at zero energy, that is, negative energy bound states emerge iff the coupling constant is incremented by an arbitrary amount (in terminology of [2] the system is at the coupling constant threshold).

The following integral operator appears in the Birman–Schwinger principle [2, 5]

$$L(k) := \sqrt{V_{12}} \left(-\Delta_x + k^2 \right)^{-1} \sqrt{V_{12}}. \quad (2)$$

$L(k)$ is analytic for $\operatorname{Re} k > 0$. Due to R1 one can use the integral representation and analytically continue $L(k)$ into the interior of the circle on the complex plane, which has its center at $k = 0$ and the radius $|b_2|$ [2]. The analytic continuation is denoted as $\tilde{L}(k) = \sum_n \tilde{L}_n k^n$, where $\tilde{L}_n$ are Hilbert-Schmidt operators.

Remark. In Sec. 2 in [2] (page 255) Klaus and Simon consider only finite range potentials. In this case $L(k)$ can be analytically continued into the whole complex plane. As the authors mention it in Sec. 9 the case of potentials with an exponential fall off requires only a minor change: the analytic continuation takes place in a circle $|k| < b_2$.

Under requirements R1-2 the operator $L(0) = \tilde{L}(0)$ is Hilbert-Schmidt and its maximal eigenvalue is equal to one

$$L(0)\phi_0 = \phi_0. \quad (3)$$

$L(0)$ is positivity-preserving, hence, the maximal eigenvalue is non-degenerate and $\phi_0 \geq 0$. We choose the normalization, where $\|\phi_0\| = 1$.

By the standard Kato–Rellich perturbation theory [5, 6] there exists $\rho > 0$ such that for $|k| \leq \rho$

$$\tilde{L}(k)\phi(k) = \mu(k)\phi(k), \quad (4)$$

where $\mu(k), \phi(k)$ are analytic, $\mu(0) = 1$, $\phi(0) = \phi_0$ and the eigenvalue $\mu(k)$ is non-degenerate. By Theorem 2.2 in [2]

$$\mu(k) = 1 - ak + O(k^2), \quad (5)$$

where

$$a = (\phi_0, (V_{12})^{1/2})^2 / (4\pi) > 0. \quad (6)$$

The orthonormal projection operators

$$\mathbb{P}(k) := (\phi(k), \cdot)\phi(k) = (\phi_0, \cdot)\phi_0 + \mathcal{O}(k), \quad (7)$$

$$\mathbb{Q}(k) := 1 - \mathbb{P}(k) \quad (8)$$

are analytic for $|k| < \rho$ as well. Our aim is to analyze the following operator function on $k \in (0, \infty)$

$$W(k) = [1 - L(k)]^{-1}. \quad (9)$$

By the Birman–Schwinger principle $\|L(k)\| < 1$ for $k > 0$, which makes $W(k)$ well-defined.

Lemma 1. *There exists $\rho_0 > 0$ such that for $k \in (0, \rho_0)$*

$$W(k) = \frac{\mathbb{P}_0}{ak} + \mathcal{Z}(k), \quad (10)$$

where $\mathbb{P}_0 := (\phi_0, \cdot)\phi_0$ and $\sup_{k \in (0, \rho_0)} \|\mathcal{Z}(k)\| < \infty$.

Proof. $\tilde{L}(k) = L(k)$ when $k \in (0, \rho)$. We get from (9)

$$W(k) = [1 - L(k)]^{-1} = [1 - L(k)]^{-1}\mathbb{P}(k) + [1 - L(k)]^{-1}\mathbb{Q}(k) \quad (11)$$

$$= [1 - \mu(k)\mathbb{P}(k)]^{-1}\mathbb{P}(k) + [1 - \mathbb{Q}(k)L(k)]^{-1}\mathbb{Q}(k) \quad (12)$$

$$= \frac{1}{1 - \mu(k)}\mathbb{P}(k) + \tilde{\mathcal{Z}}(k), \quad (13)$$

where

$$\tilde{\mathcal{Z}}(k) := [1 - \mathbb{Q}(k)L(k)]^{-1}\mathbb{Q}(k). \quad (14)$$

Note that $\sup_{k \in (0, \rho)} \|\mathbb{Q}(k)L(k)\| < 1$ because the eigenvalue $\mu(k)$ remains isolated in this range. Thus $\tilde{\mathcal{Z}}(k) = O(1)$. Using (5), (6) and (7) proves the lemma. $\square$

Remark. The singularity of $W(k)$ near $k = 0$ has been analyzed in [3] (Lemma 2.2 in [3]), see also [4]. The decomposition (10) differs in the sense that $\mathcal{Z}(k)$ is uniformly bounded in the vicinity of $k = 0$. The price we have to pay for that is the requirement R1 on the exponential fall off of V_{12} .

III. MAIN RESULT

Let us consider the Schrödinger operator for three particles in $\mathbb{R}^3$

$$H = H_0 - V_{12}(r_1 - r_2) - V_{13}(r_1 - r_3) - V_{23}(r_2 - r_3), \quad (15)$$

where r_i are particle position vectors and H_0 is the kinetic energy operator with the center of mass removed. Apart from R1-2 we shall need the following additional requirement

R3 $V_{13}, V_{23} \in L^2(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$ and $V_{13}, V_{23} \geq 0$ and $V_{23} \neq 0$.

Here we shall prove

Theorem 1. *Suppose H defined in (15) satisfies R1-3. Suppose additionally that $H \geq 0$ and $H\psi_0 = 0$, where $\psi_0 \in D(H_0)$. Then $\psi_0 = 0$.*

We defer the proof to the end of this section. Our next aim would be to derive the inequality (33)-(35).

Like in [1] we use the Jacobi coordinates $x = [\sqrt{2\mu_{12}/\hbar}](r_2 - r_1)$ and $y = [\sqrt{2M_{12}/\hbar}](r_3 - m_1/(m_1 + m_2)r_1 - m_2/(m_1 + m_2)r_2)$, where $\mu_{ij} = m_i m_j / (m_i + m_j)$ and $M_{ij} = (m_i + m_j)m_l / (m_i + m_j + m_l)$ are reduced masses (the indices i, j, l are all different). The full set of coordinates in $\mathbb{R}^6$ is labeled by ξ . In the Jacobi coordinates the kinetic energy operator takes the form

$$H_0 = -\Delta_x - \Delta_y. \quad (16)$$

Following the notation in [1] $\mathcal{F}_{12}$ denotes the partial Fourier transform in $L^2(\mathbb{R}^6)$

$$\hat{f}(x, p_y) = \mathcal{F}_{12}f(x, y) = \frac{1}{(2\pi)^{3/2}} \int d^3y e^{-ip_y \cdot y} f(x, y). \quad (17)$$

(Here and always a hat over a function denotes its Fourier transform in y -coordinates). We shall need the following trivial technical lemmas.

Lemma 2. *Suppose an operator A is positivity preserving and $\|A\| < 1$. Then $(1 - A)^{-1}$ is bounded and positivity preserving.*

Proof. A simple expansion of $(1 - A)^{-1}$ into von Neumann series. □

Lemma 3. *Suppose $g(y) \in L^2 \cap L^1(\mathbb{R}^3)$ and $g(y) \geq 0, g \neq 0$. Then*

$$\lim_{z \rightarrow +0} \int_{|p_y| \leq \epsilon_0} d^3p_y \frac{|\hat{g}|^2}{(p_y^2 + z^2)^{3/2}} = \infty \quad (18)$$

for all $\epsilon_0 > 0$.

Proof. Let us set

$$J_\epsilon(z) = \int_{|p_y| < \epsilon} d^3p_y \frac{1}{(p_y^2 + z^2)^{3/2}} \left| \int d^3y e^{ip_y \cdot y} g(y) \right|^2. \quad (19)$$

We have

$$J_\epsilon(z) \geq \int_{|p_y| < \epsilon} d^3p_y \frac{1}{(p_y^2 + z^2)^{3/2}} \left| \int d^3y g(y) \cos(p_y \cdot y) \right|^2. \quad (20)$$

Let us fix r so that

$$\int_{|y| > r} d^3y g(y) = \frac{1}{4} \|g\|_1 \quad (21)$$

Setting $\epsilon = \min[\epsilon_0, \pi/(3r)]$ we get

$$\cos(p_y \cdot y) \geq \frac{1}{2} \quad \text{if } |p_y| \leq \epsilon, |y| \leq r. \quad (22)$$

Substituting (22) and (21) into (20) we get

$$J_\epsilon(z) \geq \frac{\|g\|_1^2}{64} \int_{|p_y| < \epsilon} d^3 p_y \frac{1}{(p_y^2 + z^2)^{3/2}}. \quad (23)$$

The integral in (23) logarithmically diverges for $z \rightarrow +0$. $\square$

Remark. Lemma 2 may hold for $g(y) \in L^2(\mathbb{R}^3)$ but we could not prove this.

So let us assume that there is a bound state $\psi_0 \in D(H_0)$ at zero energy, where $\psi_0 > 0$ because it is the ground state [5]. Then we would have

$$H_0 \psi_0 = V_{12} \psi_0 + V_{13} \psi_0 + V_{23} \psi_0, \quad (24)$$

Adding the term $z^2 \psi_0$ (where here and further $z > 0$) and acting with an inverse operator on both sides of (24) gives

$$\psi_0 = [H_0 + z^2]^{-1} V_{12} \psi_0 + [H_0 + z^2]^{-1} V_{13} \psi_0 + [H_0 + z^2]^{-1} V_{23} \psi_0 \quad (25)$$

$$+ z^2 [H_0 + z^2]^{-1} \psi_0. \quad (26)$$

From now we let z vary in the interval $(0, \rho_0/2)$, where ρ_0 was defined in Lemma 1. Because the operator $[H_0 + z^2]^{-1}$ is positivity preserving [5] we obtain the inequality

$$\psi_0 \geq [H_0 + z^2]^{-1} \sqrt{V_{12}} (\sqrt{V_{12}} \psi_0) \quad (27)$$

Now let us focus on the term $\sqrt{V_{12}} \psi_0$. Using (25) we get

$$\left[1 - \sqrt{V_{12}} (H_0 + z^2)^{-1} \sqrt{V_{12}} \right] \sqrt{V_{12}} \psi_0 = \sqrt{V_{12}} [H_0 + z^2]^{-1} V_{13} \psi_0 \quad (28)$$

$$+ \sqrt{V_{12}} [H_0 + z^2]^{-1} V_{23} \psi_0 + z^2 \sqrt{V_{12}} [H_0 + z^2]^{-1} \psi_0 \quad (29)$$

And by Lemma 2

$$\sqrt{V_{12}} \psi_0 \geq \left[1 - \sqrt{V_{12}} (H_0 + z^2)^{-1} \sqrt{V_{12}} \right]^{-1} \sqrt{V_{12}} [H_0 + z^2]^{-1} V_{23} \psi_0. \quad (30)$$

It is technically convenient to cut off the wave function ψ_0 by introducing

$$\psi_1 := \psi_0(\xi) \chi_{\{|\xi| \leq b\}}, \quad (31)$$

where clearly $\psi_1 \in L^2 \cap L^1(\mathbb{R}^6)$ and $b > 0$ is fixed so that $\|V_{23} \psi_1\| \neq 0$ (which is always possible since $V_{23} \neq 0$).

Applying again Lemma 2 we get out of (30)

$$\sqrt{V_{12}}\psi_0 \geq \left[1 - \sqrt{V_{12}}(H_0 + z^2)^{-1}\sqrt{V_{12}}\right]^{-1} \sqrt{V_{12}}[H_0 + 1]^{-1}V_{23}\psi_1. \quad (32)$$

Substituting (32) into (27) gives that for all $z \in (0, \rho_0/2)$

$$\psi_0 \geq f(z) \geq 0, \quad (33)$$

where

$$f(z) = [H_0 + z^2]^{-1}\sqrt{V_{12}}\left[1 - \sqrt{V_{12}}(H_0 + z^2)^{-1}\sqrt{V_{12}}\right]^{-1} \quad (34)$$

$$\times \sqrt{V_{12}}[H_0 + 1]^{-1}V_{23}\psi_1. \quad (35)$$

Our aim would be to prove that $\lim_{z \rightarrow +0} \|f(z)\| = \infty$, which would be in contradiction with (33). Let us define

$$\Phi(x, y) := [H_0 + 1]^{-1}V_{23}\psi_1, \quad (36)$$

$$g(y) := \int dx \Phi(x, y)\sqrt{V_{12}(\alpha x)}\phi_0(x), \quad (37)$$

where ϕ_0 is defined in Sec. II.

Lemma 4. $g(y) \in L^1 \cap L^2(\mathbb{R}^3)$ and $g(y) \neq 0$.

Proof. Following [2] let us denote by $G_0(\xi - \xi', 1)$ the integral kernel of $[H_0 + 1]^{-1}$. We need a rough upper bound on $G_0(\xi, 1)$. Using the formula on p. 262 in [2] we get

$$(4\pi)^3|\xi|^4 e^{|\xi|/2}G_0(\xi, 1) = \int_0^\infty t^{-3}e^{|\xi|/2}e^{-t|\xi|^2}e^{-1/(4t)}dt \quad (38)$$

$$\leq \int_0^\infty t^{-3}e^{-3/(16t)}dt = \frac{256}{9} \quad (39)$$

Hence,

$$G_0(\xi, 1) \leq \frac{4}{9\pi|\xi|^4}e^{-|\xi|/2}. \quad (40)$$

Using $\|\sqrt{V_{12}}\phi_0\|_\infty < \infty$ we get $g(y) \in L^1 \cap L^2(\mathbb{R}^3)$ if $\Phi \in L^1 \cap L^2(\mathbb{R}^6)$. Because $\Phi \in L^2(\mathbb{R}^6)$ to prove $\Phi \in L^1(\mathbb{R}^6)$ it suffices to show that $\chi_{\{|\xi| \geq 2b\}}\Phi(\xi) \in L^1(\mathbb{R}^6)$, where b was defined after Eq. (31). This follows from (40)

$$\chi_{\{|\xi| \geq 2b\}}\Phi(\xi) \leq \chi_{\{|\xi| \geq 2b\}} \int_{|\xi'| \leq b} d^6\xi' G_0(\xi - \xi', 1)V_{23}\Psi_1(\xi') \quad (41)$$

$$\leq \chi_{\{|\xi| \geq 2b\}} \frac{4}{9\pi(|\xi| - b)^4} e^{-(|\xi| - b)/2} \|V_{23}\Psi_1\|_1 \in L^1(\mathbb{R}^6) \quad (42)$$

That $g \neq 0$ follows from the inequality $\Phi(x, y) > 0$. $\square$

Applying $\mathcal{F}_{12}$ to (34)–(35) we get

$$\hat{f}(z) = [-\Delta_x + p_y^2 + z^2]^{-1} \sqrt{V_{12}} \left[1 - \sqrt{V_{12}} (-\Delta_x + p_y^2 + z^2)^{-1} \sqrt{V_{12}} \right]^{-1} \quad (43)$$

$$\sqrt{V_{12}} [-\Delta_x + p_y^2 + 1]^{-1} \widehat{V_{23} \psi_0}. \quad (44)$$

By Lemma 1 for $|p_y| < \rho_0/2$ and $z < \rho_0/2$

$$\left[1 - \sqrt{V_{12}} \left(-\Delta_x + p_y^2 + z^2 \right)^{-1} \sqrt{V_{12}} \right]^{-1} = \frac{\mathbb{P}_0}{a \sqrt{p_y^2 + z^2}} + \mathcal{Z} \left(\sqrt{p_y^2 + z^2} \right). \quad (45)$$

From now on $z \in (0, \rho_0/2)$. Notating shortly $\chi_0(p_y) := \chi_{\{|p_y| < \rho_0/2\}}$ gives us

$$\chi_0(p_y) \hat{f}(z) = \hat{f}_1(z) + \hat{f}_2(z), \quad (46)$$

where

$$\hat{f}_1(z) = \chi_0(p_y) \frac{g(p_y)}{\sqrt{p_y^2 + z^2}} [-\Delta_x + p_y^2 + z^2]^{-1} (\sqrt{V_{12}} \varphi(x)), \quad (47)$$

$$\hat{f}_2(z) = \chi_0(p_y) [-\Delta_x + p_y^2 + z^2]^{-1} \sqrt{V_{12}} \mathcal{Z} \left(\sqrt{p_y^2 + z^2} \right) \quad (48)$$

$$\sqrt{V_{12}} [-\Delta_x + p_y^2 + 1]^{-1} (\mathcal{F}_{12} V_{23} \mathcal{F}_{12}^{-1}) \hat{\psi}_0. \quad (49)$$

The next lemma follows from the results of [1].

Lemma 5. $\sup_{z \in (0, \rho_0/2)} \|f_2(z)\| < \infty$

Proof. Let us rewrite (48)–(49) in the form

$$f_2(z) = \mathcal{A}(z) \mathcal{B}(z) \mathcal{C}(z) \psi_0, \quad (50)$$

where

$$\mathcal{A}(z) = \chi_0(p_y) [-\Delta_x + p_y^2 + z^2]^{-1} \sqrt{V_{12}} [1 + t(p_y) + z], \quad (51)$$

$$\mathcal{B}(z) = \chi_0(p_y) \mathcal{Z} \left(\sqrt{p_y^2 + z^2} \right), \quad (52)$$

$$\mathcal{C}(z) = \chi_0(p_y) \sqrt{V_{12}} [-\Delta_x + p_y^2 + 1]^{-1} [1 + t(p_y) + z]^{-1} (\mathcal{F}_{12} V_{23} \mathcal{F}_{12}^{-1}), \quad (53)$$

and $t(p_y)$ is defined as in Eq. (22) in [1]. Note that by (10) $\tilde{Z} \left(\sqrt{p_y^2 + z^2} \right)$ is a difference of two operators each of which commutes with the operator of multiplication by $[1 + t(p_y) + z]$.

That $\sup_{z \in (0, \rho_0/2)} \|\mathcal{A}(z)\|, \|\mathcal{C}(z)\| < \infty$ follows directly from the proofs of Lemmas 6,8 in [1].

$\sup_{z \in (0, \rho_0/2)} \|\mathcal{B}(z)\| < \infty$ follows from Lemma 1. $\square$

The last Lemma needed for the proof of Theorem 1 is

Lemma 6. $\lim_{z \rightarrow 0} \|f_1(z)\| = \infty$.

Proof. We get

$$\|\hat{f}_1(z)\|^2 = \frac{1}{4\pi^2} \int_{|p_y| \leq \rho_0/2} dp_y \frac{|\hat{g}(p_y)|^2}{p_y^2 + z^2} \int dx \int dx' \int dx'' \frac{e^{-\sqrt{p_y^2 + z^2}|x-x'|}}{|x-x'|} \quad (54)$$

$$\times \frac{e^{-\sqrt{p_y^2 + z^2}|x-x''|}}{|x-x''|} (\sqrt{V_{12}}(\alpha x') \phi_0(x')) (\sqrt{V_{12}}(\alpha x'') \phi_0(x'')). \quad (55)$$

There are constants $R_0, C_0 > 0$ such that

$$\int d^3x' \frac{e^{-\delta|x-x'|}}{|x-x'|} \sqrt{V_{12}}(\alpha x') \phi_0(x') \geq C_0 \frac{e^{-2\delta|x|}}{|x|} \chi_{\{|x| \geq R_0\}} \quad (56)$$

for all $\delta > 0$. Indeed, the following inequality holds for all $R_0 > 0$

$$\chi_{\{|x| \geq R_0\}} \frac{e^{-\delta|x-x'|}}{|x-x'|} \chi_{\{|x'| \leq R_0\}} \geq \frac{e^{-2\delta|x|}}{2|x|} \chi_{\{|x| \geq R_0\}}. \quad (57)$$

Substituting (57) into the lhs of (56) we obtain (56), where

$$C_0 = \frac{1}{2} \int_{|x'| \leq R_0} d^3x' \sqrt{V_{12}}(\alpha x') \phi_0(x') \quad (58)$$

and one can always choose R_0 so that $C_0 > 0$. Using (56) we get

$$\|\hat{f}_1(z)\|^2 \geq c \int_{|p_y| \leq \frac{\rho_0}{2}} dp_y \frac{|\hat{g}(p_y)|^2}{(p_y^2 + z^2)^{3/2}}, \quad (59)$$

where $c > 0$ is a constant. Now the result follows from Lemmas 3, 4. $\square$

The proof of Theorem 1 is now trivial.

Proof of Theorem 1. A bound state at threshold should it exist must satisfy inequality (33) for all $z \in (0, \rho_0/2)$. Thus $\|f(z)\|$ and, hence, $\|\chi_0 \hat{f}(z)\|$ are uniformly bounded for $z \in (0, \rho_0/2)$. By (46) and Lemmas 5, 6 this leads to a contradiction. $\square$

IV. EXAMPLE

Suppose that R2 is fulfilled and let us introduce the coupling constants $\Theta, \Lambda > 0$ in the following manner

$$H(\Theta, \Lambda) = [-\Delta_x - V_{12}] - \Delta_y - \Theta V_{13} - \Lambda V_{23}. \quad (60)$$

For simplicity, let us require that $V_{ik} \geq 0$ and $V_{ik} \in C_0^\infty(\mathbb{R}^3)$. Let $\Theta_{cr}, \Lambda_{cr}$ denote the 2-body coupling constant thresholds for particle pairs 1,3 and 2,3 respectively. On one hand, using a variational argument it is easy to show that there exists $\epsilon > 0$ such that $H(\Theta, \Lambda) > 0$ if $\Theta, \Lambda \in [0, \epsilon]$ (that is in this range $H(\Theta, \Lambda)$ has neither negative energy bound states nor a zero energy resonance) [7, 8]. On the other hand, by the Efimov effect the negative spectrum of $H(\Theta_{cr}, \Lambda)$ is not empty for $\Lambda \in (0, \Lambda_{cr})$ [3, 4]. So let us fix $\Lambda = \epsilon$ and let Θ vary in the range $[\epsilon, \Theta_{cr}]$. The energy of the ground state $E_{gr}(\Theta) = \inf \sigma(H(\Theta, \epsilon))$ is a continuous function of Θ . $E_{gr}(\Theta)$ decreases monotonically at the points where $E_{gr}(\Theta) < 0$. Because $E_{gr}(\epsilon) = 0$ there must exist $\Theta_0 \in (\epsilon, \Theta_{cr})$ such that $E_{gr}(\Theta) < 0$ for $\Theta \in (\Theta_0, \Theta_{cr})$ and $E_{gr}(\Theta_0) = 0$.

Summarizing, $H(\Theta_0, \epsilon)$ is at the 3-body coupling constant threshold. By Theorem 1 $H(\Theta_0, \epsilon)$ has a zero energy resonance but not a zero energy bound state. If $\psi_{gr}(\Theta, \xi) \in L^2(\mathbb{R}^6)$ is a wave function of the ground state defined on the interval (Θ_0, Θ_{cr}) then for $\Theta \rightarrow \Theta_0 + 0$ the wave function must totally spread (see Sec. 2 in [1]). Which means that for any $R > 0$

$$\lim_{\Theta \rightarrow \Theta_0 + 0} \int_{|\xi| < R} |\psi_{gr}(\Theta, \xi)|^2 d\xi \rightarrow 0. \quad (61)$$

Note also that if the particles 1,2 would have a ground state at the energy $e_{12} < 0$ then the ground state of the 3 body system at the energy e_{12} cannot be bound, it can only be a resonance [9].

Acknowledgments

The author would like to thank Prof. Walter Greiner for the warm hospitality at FIAS.

-
- [1] D. K. Gridnev, arXiv:0912.0414v2
 - [2] M. Klaus and B. Simon, Ann. Phys. (N.Y.) **130**, 251 (1980)
 - [3] A. V. Sobolev, Commun. Math. Phys. **156**, 101 (1993)
 - [4] D. R. Yafaev, Math. USSR-Sb. **23**, 535-559 (1974); Notes of LOMI Seminars **51** (1975) (Russian)

- [5] M. Reed and B. Simon, *Methods of Modern Mathematical Physics*, vol. 4, Academic Press/New York (1978)
- [6] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag/Berlin Heidelberg (1995)
- [7] J.-M. Richard and S. Fleck, Phys. Rev. Lett. **73**, 1464 (1994)
- [8] D. K. Gridnev and J. S. Vaagen, Phys. Rev. **C61**, 054304 (2000)
- [9] M. Klaus and B. Simon, Comm. Math. Phys. **78**, 153 (1980)